

This un-edited manuscript has been accepted for publication in Biophysical Journal and is freely available on BioFast at <http://www.biophysj.org>. The final copyedited version of the paper may be found at <http://www.biophysj.org>.

Rows of ATP synthase dimers in native mitochondrial inner membranes

Nikolay Buzhynskyy¹, Pierre Sens², Valerie Prima³, James N. Sturgis³ & Simon Scheuring^{1*}

1 - Institut Curie, UMR168-CNRS, 26 Rue d'Ulm, 75248 Paris, France.

2 – ESPCI, CNRS-UMR 7083, 10 rue Vauquelin, 75231 Paris, France.

3 - UPR-9027 LISM, IBSM, CNRS, 31 Chemin Joseph Aiguier, 13402 Marseille, France.

*Correspondence should be addressed to S. Scheuring,

Email: simon.scheuring@curie.fr

Tel.: ++33-1-42346781, Fax: ++33-1-40510636.

Running title: Supramolecular organisation of ATP synthases

ABSTRACT

The ATP synthase is a nanometric rotary machine that uses a transmembrane electrochemical gradient to form ATP. The structures of most components of the ATP synthase are known, and their organisation has been elucidated. However, the supramolecular assembly of ATP synthases in biological membranes remains unknown. Here we show with sub-molecular resolution the organisation of ATP synthases in the yeast mitochondrial inner membranes (MIM). The atomic force microscopy (AFM) images we have obtained show how these molecules form dimers with characteristic 15nm distance between the axes of their rotors through stereo-specific interactions of the membrane embedded portions of their stators. A different interaction surface is responsible for the formation of rows of dimers. Such an organisation elucidates the role of the ATP-synthase in mitochondrial morphology. Some dimers have a different morphology with 10nm stalk-to-stalk distance, in line with ATP synthases that are accessible to IF₁ inhibition. Rotation torque compensation within ATP-synthase dimers stabilizes the ATP-synthase structure in particular the stator-rotor interaction.

Keywords: atomic force microscopy; ATP synthase; membrane proteins; membrane structure; mitochondrion

INTRODUCTION

The ATP synthase is the major energy transducer protein in all cells (1, 2). The structure of the ATP synthase has been well described and can be divided into a soluble F_1 part that shows ATPase activity, and a membrane associated F_0 part that uses the proton motive force to mechanically drive F_1 . F_1 forms a ~12nm diameter head comprising three α and three β subunits housing the ADP/ATP binding sites (3), and a central stalk made of subunits γ , δ and ϵ (4), that provide an elastic coupling between F_1 and F_0 . The F_0 part forms the rotary machine. In yeast it contains an oligomeric ring of 10 c subunits (5); the number of c subunits varies among species (6, 7). This ring is associated with a peripheral stator stalk that is formed by extramembraneous subunits ‘part of b' , d , F_6 , and OSCP (the structure of this part is known (8)), and the transmembrane moiety formed by subunits a , ‘two helices of b' , e , f , g , and A6L. The structure of this last part is the least well known as only a 32 Å resolution 3D envelope (in context with the entire enzyme) is available (9). The structure of bovine IF_1 inhibitor that binds F_1 has been solved isolated (10) and complexed with F_1 (11). A large body of indirect evidence has accumulated that the ATP synthase forms dimers and oligomers, crucial for mitochondria morphology, structure, and function (12-15). In particular, F_0 subunits e and g were identified to induce dimerisation (12, 13). The bovine regulatory protein IF_1 was also shown to be associated with F_1 dimers (10, 11, 16). Detergent solubilized ATP synthase dimers were studied by cryo-EM single particle analysis (17). However direct visualization of ATP synthase dimers and oligomers in the native membrane has remained impossible. To bring light into the issue concerning the supramolecular assembly of ATP synthase holo-enzymes, we have performed an AFM (18) study of the inter-membrane space surface of native mitochondrial inner membranes (MIM) from yeast. The AFM, due to its outstanding signal-to-noise ratio, has proven to be an invaluable tool for the study of the supramolecular assembly of membrane protein in native membranes (19-21).

RESULTS AND DISCUSSION

Characterization of the mitochondria inner membranes (MIM)

Prior to the AFM studies, MIM were characterized by SDS-PAGE analysis. The hydrophobicity of the transmembrane proteins alter the migration behaviour of the proteins in the gel compared to the expected molecular masses. Therefore, we performed liquid chromatography and mass spectrometry analyses to positively identify the protein bands. These techniques revealed the presence of many different proteins including ATP synthase, succinate dehydrogenase, cytochrome bc_1 , succinate-fumarate transporter, phosphate transporter, ATP/ADP transporter, cytochrome oxidase, NADH dehydrogenase, and TIM as among abundant proteins in the studied membranes (**Fig. 1**). Furthermore, we have assembled a list of the characteristics of the typical inner mitochondrial membrane proteome (**Table 1**) that describes the expected size and form of the various complexes that might be found in MIM.

AFM analysis

Yeast MIM were adsorbed on mica and investigated by AFM. Often we found folded membranes that were opened using the AFM tip as a nano-dissector (22, 23) (**Fig. 2A**). These membranes are expected to originate from mitochondrial cristae, because of their strong curvature. After opening of the folded membrane, the AFM tip can image the protein surfaces of the mitochondrial inter-membrane space (**Fig. 2B**). Section analysis of low-resolution

topographs showed that these membranes had a thickness of $\sim 17\text{nm}$, remaining locally curved. The measured membrane thickness of $\sim 17\text{nm}$ corresponds well to the full height of ATP synthases (5). In these membranes densely packed circular complexes with a diameter of $\sim 8\text{nm}$ and inter-complex distances of $\sim 15\text{nm}$, were observed (**Fig. 2B, C, D, E**), corresponding well to the observed inter-complex distance of F_1 heads ($13.6\text{nm} - 18\text{nm}$) (24).

At first sight, the circular complexes appeared to be arranged in a hexagonal close packing (**Fig. 2B, C, D**), however upon closer examination rows of dimers were detected (**Fig. 2E**). These rows are formed by dimers at an angle of $\sim 30^\circ$ with respect to the row long axis (**Fig. 2E, 3A**). Among the typical inner mitochondrial membrane proteome, only the ATP synthase is expected to have a circular transmembrane domain, the *c*-ring (**Table 1**). Furthermore, the ring dimensions correlate well with the size of the *c*-ring (6, 7, 25), the inter-ring distances correspond well to the known ATP synthase neighbour spacing in yeast mitochondria (24), and the membrane thickness fits well the total height of the entire ATP synthase (5, 9). Therefore, based on the characteristic size and shape of the observed circular complexes and their presence in rows of dimers we tentatively assign them as a view of the *c*-ring inter-membrane space surface of the ATP synthase.

High-resolution raw data images revealed protein topographies in the centre of the *c*-ring and neighbouring the *c*-ring (**Fig. 3A**). These central and peripheral proteins protrude $\sim 1\text{nm}$ further than the *c*-ring and make the rotor appear as an indentation. This may appear counter-intuitive, but is in line with structural information on the ATP synthase hollow enzyme: first, a cryo-EM analysis of the detergent-solubilized holo-enzyme monomer showed that the stator F_0 part protrudes further into the inter-membrane space than the rotor (9). This cryo-EM 3D-map (9) has proven to be accurate since the atomic structure of the extramembraneous part of the peripheral stalk matches well the EM envelope (8). Second, an EM study of detergent-solubilized ATP synthase dimers revealed a density in the centre of the *c*-ring on the inter-membrane space surface (17). While the protein located at this density remained unassigned, this feature was interpreted as F_0 bridging important for dimer formation and implicated in cristae biogenesis (17). Third, equally, the cryo-EM 3D-map of the related V-ATP synthase features a well defined density that ‘plugs’ the *c*-ring on the surface opposite to F_1 (26). Fourth, AFM analysis of purified and reconstituted chloroplast *c*-rings with associated additional subunits revealed a *c*-ring plug on one surface (27). The peripheral protrusions correspond to stator F_0 transmembrane subunits *a*, ‘two transmembrane helices of *b*’, *e*, *g* and A6L, in agreement with the cryo-EM map (9). Cross-correlation averaging showed that these peripheral proteins have three major protruding domains and have a preserved position between ATP synthases within the dimer and between adjacent dimers within the row (**Fig. 3B**). Based on our images and a strong body of structural and biochemical data (12, 13), we propose that dimerization subunits *e*, *g* and the dimerizing portions of *b* are positioned at these interfaces (**Fig. 3C**), without a precise structural assignment of these subunits to the features observed by AFM. Dimerization over subunits *e*, *g* and over subunits *b* are structurally not equivalent. Both are sufficient to induce independently dimer formation, but are together needed for oligomer formation, in line with the fact that either deletion has drastic effects on mitochondria morphology (13). We propose the *e/g*-dimerization surface is located within the dimer, while *b*-dimerization, shown to efficiently cross-link at amino-acid positions located in the intermembrane space (13), acts along the dimer row. Notably, we imaged inter-membrane space protrusions of neighbouring F_0 close to this interface (**Fig. 3B, C**).

Most enzyme dimers reveal a stalk-to-stalk distance of 15nm , however a second class of ATP synthase shows 10nm separation (**Fig. 3D**; see also **Fig. 3A, B**). We propose that these two classes represent active ATP synthase enzyme dimers and ATPase enzyme dimers that

are accessible for IF₁ inhibition (**Fig. 3E**). Two IF₁ inhibitor molecules were shown to clamp two F₁ together (11). It appears suggestive that the small molecule IF₁ requires preformed ATP synthase dimers. Inhibited dimeric F₁ are linked together between α and β subunits of each F₁. In this assembly the stalks of the two ATP synthases are closer together ($\sim 10\text{nm}$) (11) than could be achieved if two of the bulkier subunit edges of each F₁ faced each other ($\sim 13\text{nm}$). Therefore we suggest that the dimers with 10nm stalk-to-stalk distance are sterically accessible for IF₁ inhibition. The decrease of distance between the two ATP synthases requires that the F_O stator moieties are displaced, away from their position close to the dimer centre (**Fig. 3B**). We estimate that both ATP synthases must rotate $\sim 30^\circ$ to accomplish this novel dimer conformation, breaking the intramembrane contacts between *e/g* dimerization surfaces (**Fig. 3B, E right**). We furthermore observe that this conformation appears incompetent for oligomer formation in rows, i.e. the dimerization surface of *b* subunits are not correctly exposed to neighbouring ATP synthases any longer, and thus they do not participate in the oligomers (**Fig. 3A**).

Theoretical considerations on the functional role of ATP synthase dimers

In the ATP synthase, the energy from either proton translocation or ATP-hydrolysis is converted into rotary motion. Both processes inject fixed amounts of energy (depending on the ATP/ADP concentration and/or the transmembrane proton potential) into the system, which is converted into mechanical torque on the moving parts of the machine. Because no net force can be produced by the activity of the enzyme, equal but opposite torques τ_A and τ_B must be generated on the rotor and the stator at the active site (**Fig. 4A**).

The energy influx per unit time $d_t E$ is dissipated both by internal friction within the protein and by viscous friction with the membrane environment. Conservation of energy imposes that the energy influx is equal to the total dissipation:

$$d_t E = P_A + P_B + P_i = \frac{1}{2} \zeta_A \Omega_A^2 + \frac{1}{2} \zeta_B \Omega_B^2 + \frac{1}{2} (\zeta_i + \zeta_{AB}) (\Omega_A + \Omega_B)^2 \quad (1)$$

where the dissipation has been split into P_A and P_B that depend on the individual motion of A and B with respect to the membrane, and P_i that depends on the relative motion of A with respect to B. The latter includes the membrane flow in the vicinity of the A – B interface and the energy converted into internal friction within the protein, both at the proton translocation site (transmembrane interface between rotor and stator) and at the ADP/ATP conversion site (interface between the stator F₁(α , β) subunits and the rotor γ subunit). The internal friction at these sites includes the energy dissipated by pumping protons or synthesizing ATP. Each type of motion is characterized by a friction coefficient ζ , which relates the dissipation to the angular velocities Ω_A and Ω_B of the rotor and the stator.

The torque of A and B can be written $\tau_A = \partial(d_t E)/\partial\Omega_A$ and $\tau_B = \partial(d_t E)/\partial\Omega_B$. Torque balance (equal but opposite torques τ_A and τ_B) determines the velocities of the rotor and stator:

$$\Omega_B = \frac{\zeta_A}{\zeta_B} \Omega_A \quad \Omega_A + \Omega_B = \frac{\tau}{\zeta_i + \zeta_{AB} + \frac{\zeta_A \zeta_B}{\zeta_A + \zeta_B}} \quad (2)$$

In two dimensions, the hydrodynamic friction force is expected to be proportional to the membrane viscosity and the velocity of the object $v = \Omega R$ (within logarithmic corrections, $R \approx 10\text{nm}$ the distance from the axis of rotation), giving $\tau_A \approx \tau_B \approx \mu_m R^2 \approx 10^{-25} \text{J.s}$, where

$\mu_m \approx 10^{-9} J.s/m^2$ is the membrane viscosity (28). Therefore, we expect the stator to actually move at about the same angular velocity as the rotor for an isolated ATP synthase monomer. The hydrodynamic friction due to the motion of A with respect to B depends on the gradient of the velocity field near the A–B interface, and is of order $\zeta_{AB} \approx (R/a)\zeta_A \approx 10^{-24} J.s$, where $a \approx 1nm$ characterizes the distance between rotor and stator. The torque produced by ATP hydrolysis is of order $\tau \approx 50 pN.nm$ (29), thus if the friction was solely due to hydrodynamics, one would expect an angular velocity of order 10,000 cycles per second (Hz). The maximal velocity of the F1- γ part of the protein in solution is about 50 times smaller (250 Hz) (30), which shows that the internal friction dominates the hydrodynamic drag, and must be of order $\zeta_i \approx 3.10^{-23} J.s$. The hydrodynamic drag in the membrane is thus much smaller than the internal friction and does not reduce the efficiency of the enzyme. This value correlates well with measurements on the angular velocity of F1-ATPase with attached actin filament as a function of the filament length (30). Similarly to Eq (2), the angular velocity of the actin-bound rotor is given by $\Omega = \tau / (\zeta_i + \zeta_{actin})$, where the friction due to the actin $\zeta_{actin} \approx \eta l_{actin}^3$ depends on the actin length l_{actin} and the water viscosity η (28). The velocity of the actin-free enzyme (250Hz) drops 20 fold to 12Hz when a 1 μm actin filament is attached (30). This leads to the independent estimate $\zeta_i \approx \zeta_{actin} / 20 \approx 5.10^{-23} J.s$.

Our observations show that ATP synthases are functional dimers, and provide evidence that dimers undergo a conformational change between the ATP-producing and ATP-consuming state. One may wonder whether rotating objects working in close contact might influence each other through hydrodynamic flow in the membrane. Hydrodynamic forces can be calculated from the viscous stress per unit protein length $\mu_m \nabla v$ (the membrane viscosity multiplied by the gradient of the velocity field). The hydrodynamic force shown (**Fig.4B, blue arrow**) for the ATP-producing state should stabilize the rotor stator interactions. Nevertheless, its magnitude is quite small (of order $f \approx \mu_m R^2 \Omega \approx 0.1 pN$) and should not have a strong influence on the protein conformation. On the other hand, the torque resulting from the enzyme activity is quite large and leads to forces of order $f \approx \tau / R \approx 5 pN$ on the stator (**Fig.4B, red arrows**). These forces stabilize the stator/stator interactions within the dimer in the ATP-producing state, and destabilize them in the ATP-consuming state. They reduce the energy gain from stator-stator interactions by an amount $\delta E \approx \tau a / R \approx 2k_B T$. This is a quite sizeable effect, as these interactions are expected to be fairly small, thus facilitating a conformational change crucial to the inhibition of the enzyme in the ATP consumption state.

It is worth noting that the lack of viscous losses and the efficiency of the machine rely on its nanometric size. If the machine were ten times larger (producing 30 ATP/cycle), the hydrodynamic drag (which scales like R^2) would dominate the internal friction and the enzyme's efficiency would be appreciably reduced.

CONCLUSIONS

Since single molecule measurements have shown that F₁ rotates the central stalk counter-clockwise upon ATP hydrolysis (29), it is reasonable to assume that the rotation is clockwise during ATP synthesis. In contrast to a previous report (31), the torque force is generated at the interface between the transmembrane portions of the rotor and the stator. As a consequence, while the rotor rotates clockwise the stator, in reaction, rotates counter-clockwise around it in a fluid membrane where nothing holds the entire machine in place. This will make the ATP synthase as a whole rotate around an axis approximately at its centre of mass and move in the membrane. Such movements dissipate energy. In the oligomeric

assembly in contrast, the torques on the stators of each molecule are compensated and only the rotors will move. Is the blocking of energy dissipation through movement of ATP synthases in the membrane sufficient to explain the increased activity of oligomeric versus monomeric ATP synthases (15)? Apparently not, the major source of energy dissipation lies inside the ATP synthase, driving ATP synthesis, and the energy provided by proton translocation would produce F_0 rotation ~ 40 times faster than observed (32) if it was only opposed to the viscous friction in the membrane as discussed above. The energy dissipation through viscous drag is thus negligible compared to friction at both rotor-stator interfaces inside the ATP synthase. Therefore we conclude that oligomer formation has rather a structural role by assuring the integrity of the ATP synthase structure, particularly by stabilizing the weak F_0 rotor-stator interfaces. Notably, the rotor movement of each ATP synthase will drive the stator of the other towards its rotor. Taking together, the torque during ATP synthesis stabilizes the interfaces within each ATP synthase, within dimers and oligomers, and as a consequence cristae morphology. In contrast ATP hydrolysis destabilizes dimers by pulling the stators apart, allowing the monomers to approach each other and favour F_1 binding. Indeed, in this conformation the rotors come close together (10nm) and may hinder each other's rotation at their interface (**Fig. 3E, right**).

We observed, using high-resolution AFM, rows of ATP synthase dimers in yeasts MIM, thus providing first direct evidence of the organization of ATP synthase in dimers and oligomers in native membranes. The proposed model describes the interactions within ATP synthase dimers and evidences their functional role. While atomic structures strongly advanced our knowledge on individual proteins, AFM studies of native membranes visualize the supramolecular assembly of multiple membrane proteins in their native environment (19-21) and provide complementary data for a complete structural description. AFM images the supercomplexes into which structures can be docked (33), and providing insight into the function of these integrated systems.

METHODS

Membrane preparation: Mitochondria were purified from yeast (*Saccharomyces cerevisiae* strain W303) cells grown overnight under oxidative conditions in YP-glycerol media at 30°C. Mitoplasts and outer membrane fragments were prepared as described (34). Mitoplasts were broken by a single passage through a French pressure cell in 1mM Tris pH 7.5. The resulting inner mitochondrial membranes were collected and washed by centrifugation. Membranes were stored at 4° for AFM analysis.

Atomic force microscopy (AFM): Mica supports were immersed in 40- μ l adsorption buffer (10 mM Tris-HCl, pH 7.4, 150 mM KCl, 25 mM MgCl₂). Subsequently, 3 μ l of membrane solution were injected into the buffer drop. After ~ 30 min, the sample was rinsed with recording buffer (10 mM Tris-HCl, pH 7.4, 150 mM KCl). The AFM (18) was operated in contact mode at ambient temperature and pressure. Imaging was performed with a commercial Nanoscope-E AFM (Veeco, Santa Barbara, CA, USA) equipped with a 160- μ m scanner (J-scanner) and oxide-sharpened Si₃N₄ cantilevers (length 100 μ m; k = 0.09 N/m; Olympus Ltd., Tokyo, Japan). For imaging minimal loading forces of ~ 100 pN were applied, at scan frequencies of 4-7 Hz using optimized feedback parameters and manually accounting for force drift.

Data analysis: All image treatment and analysis of AFM topographs were performed using self written routines for the ImageJ image processing package (33, 35) and IGOR PRO (Wavemetrics, Portland, OR, USA).

ACKNOWLEDGEMENTS

This study was supported by the INSERM (Institut National de la Santé et Recherche Médicale) and INSERM Avenir, CNRS (Centre National de la Recherche Scientifique) and the grant (ANR-06-NANO-023-01).

REFERENCES

1. Boyer, P. 1997. The ATP synthase - a splendid molecular machine. *Annu Rev Biochem* 66:717-749.
2. Stock, D., C. Gibbons, I. Arechaga, A. G. Leslie, and J. E. Walker. 2000. The rotary mechanism of ATP synthase. *Curr. Opin. Struct. Biol.* 10:672-679.
3. Abrahams, J. P., A. G. W. Leslie, R. Lutter, and J. E. Walker. 1994. Structure at 2.8 angstrom resolution of F₁-ATPase from bovine heart mitochondria. *Nature* 370:621-628.
4. Gibbons, C., M. G. Montgomery, A. G. Leslie, and J. E. Walker. 2000. The structure of the central stalk in bovine F₁-ATPase at 2.4 Å resolution. *Nat Struct Biol* 7:1055-1061.
5. Stock, D., A. G. Leslie, and J. E. Walker. 1999. Molecular Architecture of the Rotary Motor in ATP Synthase. *Science* 286:1700-1705.
6. Seelert, H., A. Poetsch, N. A. Dencher, A. Engel, H. Stahlberg, and D. J. Müller. 2000. Proton powered turbine of a plant motor. *Nature* 405:418-419.
7. Meier, T., P. Polzer, K. Diederichs, W. Welte, and P. Dimroth. 2005. Structure of the rotor ring of F-Type Na⁺-ATPase from *Ilyobacter tartaricus*. *Science* 308:659-662.
8. Dickson, V. K., J. A. Silvester, I. M. Fearnley, A. G. Leslie, and J. E. Walker. 2006. On the structure of the stator of the mitochondrial ATP synthase. *EMBO J* 25:2911-2918.
9. Rubinstein, J. L., J. E. Walker, and R. Henderson. 2003. Structure of the mitochondrial ATP synthase by electron cryomicroscopy. *EMBO J* 22:6182-6192.
10. Cabezon, E., M. Runswick, A. Leslie, and J. Walker. 2001. The structure of bovine IF(1), the regulatory subunit of mitochondrial F-ATPase. *EMBO J* 20:6990-6996.
11. Cabezon, E., M. G. Montgomery, A. G. W. Leslie, and J. E. Walker. 2003. The structure of bovine F₁-ATPase in complex with its regulatory protein IF1. *Nat Struct Mol Biol* 10:744-750.
12. Arnold, I., K. Pfeiffer, W. Neupert, R. Stuart, and H. Schagger. 1998. Yeast mitochondrial F₁F₀-ATP synthase exists as a dimer: identification of three dimer-specific subunits. *EMBO J* 17:7170-7178.
13. Paumard, P., J. Vaillier, B. Coulary, J. Schaeffer, V. Soubannier, D. M. Mueller, D. Brèthes, J.-P. di Rago, and J. Velours. 2002. The ATP synthase is involved in generating mitochondrial cristae morphology. *EMBO J* 21:221-230.
14. Allen, R., C. Schroeder, and A. Fok. 1989. An investigation of mitochondrial inner membranes by rapid-freeze deep-etch techniques. *J. Cell. Biol.* 108:2233-2240.
15. Bornhovd, C., F. Vogel, W. Neupert, and A. S. Reichert. 2006. Mitochondrial Membrane Potential Is Dependent on the Oligomeric State of F₁F₀-ATP Synthase Supracomplexes. *J. Biol. Chem.* 281:13990-13998.
16. Cabezon, E., I. Arechaga, P. Jonathan, G. Butler, and J. Walker. 2000. Dimerization of bovine F₁-ATPase by binding the inhibitor protein, IF1. *J Biol Chem* 275:28353-28355.
17. Minauro-Sanmiguel, F., S. Wilkens, and J. J. Garcia. 2005. Structure of dimeric mitochondrial ATP synthase: Novel F₀ bridging features and the structural basis of mitochondrial cristae biogenesis. *PNAS* 102:12356-12358.
18. Binnig, G., C. Gerber, E. Stoll, T. R. Albrecht, and C. F. Quate. 1987. Atomic resolution with atomic force microscopy. *Europhys. Lett.* 3:1281-1286.
19. Fotiadis, D., Y. Liang, S. Filipek, D. A. Saperstein, A. Engel, and K. Palczewski. 2003. Atomic-force microscopy: Rhodopsin dimers in native disc membranes. *Nature* 421:127-128.
20. Scheuring, S., and J. N. Sturgis. 2005. Chromatic adaptation of photosynthetic membranes. *Science* 309:484-487.
21. Buzhynskyy, N., R. K. Hite, T. Walz, and S. Scheuring. 2007. The supramolecular architecture of junctional microdomains in native lens membranes. *EMBO Reports* 8:51-55.
22. Fotiadis, D., L. Hasler, D. J. Müller, H. Stahlberg, J. Kistler, and A. Engel. 2000. Surface tongue-and-groove contours on lens MIP facilitate cell-to-cell adherence. *J. Mol. Biol.* 300:779-789.
23. Scheuring, S., J. N. Sturgis, V. Prima, A. Bernadac, D. Lévy, and J.-L. Rigaud. 2004. Watching the photosynthetic apparatus in native membranes. *Proc. Natl. Acad. Sci. U.S.A.* 101:11293-11297.
24. Nicastro, D., A. Frangakis, D. Typke, and W. Baumeister. 2000. Cryo-electron tomography of neurospora mitochondria. *J Struct Biol* 129:48-56.
25. Stahlberg, H., D. J. Müller, K. Suda, D. Fotiadis, A. Engel, U. Matthey, T. Meier, and P. Dimroth. 2001. Bacterial ATP synthase has an undecameric rotor. *EMBO Reports* 2:229-235.
26. Gregorini, M., J. Wang, X.-S. Xie, R. A. Milligan, and A. Engel. 2007. Three-dimensional reconstruction of bovine brain V-ATPase by cryo-electron microscopy and single particle analysis. *J Struct Biol* In Press.
27. Seelert, H., N. A. Dencher, and D. J. Müller. 2003. Fourteen Protomers Compose the Oligomer III of the Proton-rotor in Spinach Chloroplast ATP Synthase. *Journal of Molecular Biology* 333:337-344.
28. Happel, J., and H. Brenner. 1991. *Low Reynolds Number Hydrodynamics*. Kluwer, Dordrecht.

29. Noji, H., R. Yasuda, M. Yoshida, and K. Kinosita. 1997. Direct observation of the rotation of F₁-ATPase. *Nature* 386:299-302.
30. Yasuda, R., H. Noji, K. Kinosita, and M. Yoshida. 1998. F₁-ATPase Is a Highly Efficient Molecular Motor that Rotates with Discrete 120° Steps. *Cell* 93:1117-1124.
31. Rexroth, S., J. Meyer zu Tittingdorf, H. Schwamann, F. Krause, H. Seelert, and N. Dencher. 2004. Dimeric H⁺-ATP synthase in the chloroplast of *Chlamydomonas reinhardtii*. *Biochim Biophys Acta* 1658:202-211.
32. Ueno, H., T. Suzuki, K. Kinosita Jr., and M. Yoshida. 2005. ATP-driven stepwise rotation of FoF₁-ATP synthase. *Proc Natl Acad Sci* 102:1333-1338; .
33. Scheuring, S., T. Boudier, and J. N. Sturgis. 2007. From high-resolution AFM topographs to atomic models of supramolecular assemblies. *J. Struct. Biol.* accepted.
34. Mannella, C. 1982. Structure of the outer mitochondrial membrane: ordered arrays of porelike subunits in outer-membrane fractions from *Neurospora crassa* mitochondria. *J Cell Biol* 94:680-687.
35. Rasband, W. S. 1997-2005. ImageJ. U.S. National Institutes of Health, Bethesda, Maryland, USA, <http://rsb.info.nih.gov/ij/>.
36. Grigorieff, N. 1998. Three-dimensional structure of bovine NADH:ubiquinone oxidoreductase (complex I) at 22 Å in ice. *J Mol Biol* 277:1033-1046.
37. Abdrakhmanova, A., V. Zickermann, M. Bostina, M. Radermacher, H. Schagger, S. Kerscher, and U. Brandt. 2004. Subunit composition of mitochondrial complex I from the yeast *Yarrowia lipolytica*. *Biochim Biophys Acta* 1658:148-156.
38. Sazanov, L., and P. Hinchliffe. 2006. Structure of the Hydrophilic Domain of Respiratory Complex I from *Thermus thermophilus*. *Science* 311:1430-1436.
39. Sun, F., X. Huo, Y. Zhai, A. Wang, J. Xu, D. Su, M. Bartlam, and Z. Rao. 2005. Crystal structure of mitochondrial respiratory membrane protein complex II. *Cell* 121:1043-1057
40. Jormakka, M., S. Tornroth, B. Byrne, and S. Iwata. 2002. Molecular basis of proton motive force generation: structure of formate dehydrogenase-N. *Science* 295:1863-1868.
41. Yankovskaya, V., R. Horsefield, S. Tornroth, C. Luna-Chavez, H. Miyoshi, C. Leger, B. Byrne, G. Cecchini, and S. Iwata. 2003. Architecture of succinate dehydrogenase and reactive oxygen species generation. *Science* 299:700-704.
42. Xia, D., C. A. Yu, H. Kim, J. Z. Xia, A. M. Kachurin, L. Zhang, L. Yu, and J. Deisenhofer. 1997. Crystal structure of the cytochrome bc₁ complex from bovine heart mitochondria. *Science* 277:60-66.
43. Iwata, S., J. W. Lee, K. Okada, J. K. Lee, M. Iwata, B. Rasmussen, T. A. Link, S. Ramaswamy, and B. K. Jap. 1998. Complete structure of the 11-subunit bovine mitochondrial cytochrome bc₁ complex. *Science* 281:64-67.
44. Hunte, C., J. Koepke, C. Lange, T. Rossmann, and H. Michel. 2000. Structure at 2.3 Å resolution of the cytochrome bc₁(1) complex from the yeast *Saccharomyces cerevisiae* co-crystallized with an antibody Fv fragment. *Structure* 8:669-684.
45. Tsukihara, T., H. Aoyama, E. Yamashita, T. Tomizaki, H. Yamaguchi, K. Shinzawa-Itoh, R. Nakashima, R. Yaono, and S. Yoshikawa. 1996. The whole structure of the 13-subunit oxidized cytochrome c oxidase at 2.8 Å. *Science* 272:1136-1144.
46. Pebay-Peyroula, E., C. Dahout-Gonzalez, R. Kahn, V. Trezeguet, G. Lauquin, and G. Brandolin. 2003. Structure of mitochondrial ADP/ATP carrier in complex with carboxyatractyloside. *Nature* 426:39-44.
47. Bauer, M., S. Hofmann, W. Neupert, and M. Brunner. 2000. Protein translocation into mitochondria: the role of TIM complexes. *Trends Cell Biol* 10:25-31.
48. Koehler, C. M. 2004. NEW DEVELOPMENTS IN MITOCHONDRIAL ASSEMBLY. *Annual Review of Cell and Developmental Biology* 20:309-335.
49. Pfanner, N., and A. Geissler. 2001. Versatility of the mitochondrial protein import machinery. *Nat Rev Mol Cell Biol* 2:339-349.
50. Reif, S., O. Randelj, G. Domanska, E. Dian, T. Krimmer, C. Motz, and J. Rassow. 2005. Conserved mechanism of Oxal insertion into the mitochondrial inner membrane. *J Mol Biol* 354:520-528.

Table 1: Inner mitochondrial membrane proteins:

	kDa	# Subunits ¹	# TMH ²	Soluble domain inter ³ /matrix height (Å)	Biological unit	TMD ⁴ dimensions	References
(Complex I) NADH:ubiquinone oxidoreductase	~ 980	46 ⁶	~ 50	matrix (140)	monomer	200Å x 80Å	(36), (37), (38)
(Complex II) Succinate:ubiquinone oxidoreductase	~ 130	4	6	matrix (~80)	monomer ⁶ trimer ⁷	45Å x 45Å (monomer) 80Å x 80Å (trimer)	(39), (40), (41)
(Complex III) Ubiquinone:cytochrome <i>c</i> oxidoreductase Cytochrome <i>bc</i> ₁ complex	~ 250	7	13	matrix (~75)	dimer	70Å X 90Å	(42), (43), (44)
(Complex IV) Cytochrome <i>c</i> oxidase	~ 200	12 ⁵ 13 ⁶	28	matrix inter (~20)	dimer ⁸	200Å x 100Å	(45)
(Complex V) ATP synthase	~ 600	21	20 (<i>c</i> -ring) ~ 11 ⁹ (F _O -stator)	matrix inter (~130) (~20)	monomer ¹ dimer ⁸	60Å circular (<i>c</i> -ring) 50Å x 70Å (F _O -stator)	(5), (11), (4), (8), (9)
(Mitochondrial carriers: 35 members) ATP/ADP transporter ¹⁰	~ 33	1	6	matrix inter (~10) (~10)	monomer dimer ⁸	40Å x 40Å	(46)
TIM 22 complex	~ 300	5	17				(47), (48)
TIM22	~ 22	1 (4x)	4	matrix	tetramer	-	
TIM54	~ 54	1	1-2				
TIM 23 complex	~ 340	8	16 ⁷				(47), (49)
TIM23	~ 23	1 (2x)	4	matrix	dimer	-	
TIM17	~ 17	1 (2x)	4		dimer	-	
TIM44	~ 60	1 (2x)		matrix	dimer	-	
Hsp70	~ 70	1 (2x)		matrix	dimer	-	
Oxa1	~ 450	1	5	-	monomer	-	(50)

¹ Based on X-ray or EM structures or biochemical analysis (see references)

² TMH = Transmembrane helices ; Based on X-ray or EM structures or sequence based structure prediction (see references)

³ inter = Intermembrane space ; Based on X-ray or EM structures or sequence based structure prediction (see references)

⁴ TMD = Transmembrane domain ; Based on X-ray or EM structures or sequence based structure prediction (see references)

⁵ Yeast mitochondria

⁶ Mammalian mitochondria

⁷ Bacterial homologue

⁸ Evidenced to participate in supercomplex formation in different stoichiometry

⁹ Comprising subunits 6, 8, *f*, *i*, 4, *g*

¹⁰ representative members of the mitochondrial carrier family

FIGURE LEGENDS

Fig.1) Coomassie brilliant blue stained SDS-PAGE of yeast mitochondrial inner membrane fractions.

Bands labelled and marked with solid lines were identified on the basis of liquid chromatography and mass spectrometry analysis and gel migration behaviour. Bands marked with dotted lines were not identified.

Fig. 2) Overview AFM analysis of mitochondrial inner membranes (MIM).

- A) MIM adsorbed on mica. The folded parts of the membrane are marked with white arrows, the membrane area opened with the AFM tip is marked by the red arrow.
- B) Topograph of the flattened membrane area, shown in A). The AFM tip contours circular complexes of ~ 8 nm in diameter and inter-complex distances of ~ 15 nm corresponding to *c*-rings of ATP synthase.
- C) and D) Overview images reveal arrays of *c*-ring complexes densely packed.
- E) Row of intercalated dimers. The *c*-ring rotor appears as a ~ 1 nm indentation.

Fig. 3) The supramolecular architecture of ATP synthases.

- A) High resolution AFM topograph of mitochondrial inner membrane. Right: ATP synthase dimers are outlined.
- B) Average topographies. Left: Dimer with 15nm stalk-to-stalk distance. Right: Dimer with 10nm stalk-to-stalk distance. Bottom: Outlines indicate the molecular packing along a row of dimers.
- C) Schematic representation of dimer and oligomer assembly. The positions of the molecules correspond to the average topograph (B bottom panel). The assignment of subunits *e*, *g* at dimer interfaces, and transmembrane portion of *b* at interfaces along the dimer row, are based on (12, 13). Left: black arrows indicate rotor and stator rotation torque direction.
- D) Stalk-to-stalk distance histogram ($n = 58$). Two populations of dimers are distinguished, most dimers have 15nm stalk-to-stalk distance (white arrows in A) a minor fraction has 10nm stalk-to-stalk distance (yellow arrows in A).
- E) Schematic representation of active (left) and IF₁ inhibited (right) ATP synthase dimers. The rotation torques stabilize the dimer during ATP synthesis and separate F_O stator portions during ATP hydrolysis.

Fig. 4) Theoretical considerations about the ATP synthase dimer formation.

- A) Monomer of ATP-synthase; opposite torques τ_A and τ_B are generated on the rotor A and stator B.
- B) Functional dimers of ATP synthase. Hydrodynamic (blue arrows) and torque (red arrows) forces stabilize the interaction within the ATP-synthase dimer during the ATP production; while during the consumption the forces tend to destabilize the dimer.




Figure 1




Figure 2




Figure 4